

CODE

0

PAPER-1

P1-14-0

1205510

Time : 3 Hours

Maximum Marks : 180

Please read the instructions carefully. You are allotted 5 minutes specifically for this purpose.

INSTRUCTIONS

A. General

1. This booklet is your Question Paper. Do not break the seal of this booklet before being instructed to do so by the invigilators.
2. The question paper CODE is printed on the left hand top corner of this sheet and on the back cover page of this booklet.
3. Blank spaces and blank pages are provided in the question paper for your rough work. No additional sheets will be provided for rough work.
4. Blank papers, clipboards, log tables, slide rules, calculators, cameras, cellular phones, pagers and electronic gadget of any kind are NOT allowed inside the examination hall.
5. Write your Name and Roll number in the space provided on the back cover of this booklet.
6. Answers to the questions and personal details are to be filled on an Optical Response Sheet, which is provided separately. The ORS is a doublet of two sheets - upper and lower, having identical layout. The upper sheet is a machine-gradable Objective Response Sheet (ORS) which will be collected by the invigilator at the end of the examination. The upper sheet is designed in such a way that darkening the bubble with a ball point pen will leave an identical impression at the corresponding place on the lower sheet. You will be allowed to take away the lower sheet at the end of the examination. (see Figure-1 on the back cover page for the correct way of darkening the bubbles for valid answers).
7. Use a black ball point pen only to darken the bubbles on the upper original sheet. Apply sufficient pressure so that the impression is created on the lower sheet. See Figure-1 on the back cover page for appropriate way of darkening the bubbles for valid answers.
8. DO NOT TAMPER WITH / MUTILATE THE ORS OR THIS BOOKLET.
9. On breaking the seal of the booklet check that it contains 28 pages and all the 60 questions and corresponding answer choices are legible. Read carefully the instruction printed at the beginning of each section.

B. Filling the right part of the ORS

10. The ORS also has a CODE printed on its left and right parts.
11. Verify that the CODE printed on the ORS (on both the left and right parts) is the same as that on this booklet and put your signature in the Box designated as R4.
12. IF THE CODES DO NOT MATCH, ASK FOR A CHANGE OF THE BOOKLET / ORS AS APPLICABLE.
13. Write your Name, Roll No. and the name of centre and sign with pen in the boxes provided on the upper sheet of ORS. Do not write any of this anywhere else. Darken the appropriate bubble UNDER each digit of your Roll No. in such way that the impression is created on the bottom sheet. (see example in Figure 2 on the back cover)

C. Question Paper Format

- The question paper consists of three parts (Physics, Chemistry and Mathematics). Each part consists of two sections.
14. Section 1 contains 10 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which ONE OR MORE THAN ONE are correct.
 15. Section 2 contains 10 questions. The answer to each of the questions is a single-digit integer, ranging from 0 to 9 (both inclusive).

DO NOT BREAK THE SEAL WITHOUT BEING INSTRUCTED TO DO SO BY THE INVIGILATOR



Please read the last page of this booklet for rest of the instructions.

PART I : PHYSICS

PHYSICS

SECTION - 1 : (One or More Than One Options Correct Type)

This section contains 10 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which ONE or MORE THAN ONE are correct.

1. Let $E_1(r)$, $E_2(r)$ and $E_3(r)$ be the respective electric fields at a distance r from a point charge Q , an infinitely long wire with constant linear charge density λ , and an infinite plane with uniform surface charge density σ . If $E_1(r_0) = E_2(r_0) = E_3(r_0)$ at a given distance r_0 , then

(A) $Q = 4\sigma\pi r_0^2$

(B) $r_0 = \frac{\lambda}{2\pi\sigma}$

(C) $E_1(r_0/2) = 2E_2(r_0/2)$

(D) $E_2(r_0/2) = 4E_3(r_0/2)$

2. Heater of an electric kettle is made of a wire of length L and diameter d . It takes 4 minutes to raise the temperature of 0.5 kg water by 40 K. This heater is replaced by a new heater having two wires of the same material, each of length L and diameter $2d$. The way these wires are connected is given in the options. How much time in minutes will it take to raise the temperature of the same amount of water by 40 K?

(A) 4 if wires are in parallel

(B) 2 if wires are in series

(C) 1 if wires are in series

(D) 0.5 if wires are in parallel

Space for Rough Work

Handwritten calculations for Question 1:

$$\frac{1}{4\pi\epsilon_0} \frac{Q}{r_0^2} = \frac{\sigma}{2\epsilon_0} \quad \sqrt{\frac{Q}{4\pi\sigma}} = r_0 \quad = \frac{1}{2\pi\epsilon_0} \frac{\lambda}{r_0} = \frac{\sigma}{\epsilon_0}$$

$$Q = 4\pi\sigma r_0^2 \quad \frac{\lambda}{r_0} = \epsilon_0 \sigma$$

Handwritten calculations for Question 2:

$$\frac{5 \times 60}{4\pi d^2} = 0.5 C \times 40 \times \frac{1}{4\pi\epsilon_0} \frac{Q \times 4\pi \times 10^{-2} \times 4}{\lambda^2}$$

$$\frac{5 \times 60}{4\pi \epsilon_0} = 0.5 \times 40 \times \frac{Q \times 4\pi}{\lambda^2}$$

$$\frac{1}{4\pi\epsilon_0} \frac{Q \times 4}{r_0^2} = \frac{1}{4\pi\epsilon_0} \frac{Q \times 4}{\sigma}$$

$$\frac{PL \times 6}{4\pi d^2} = \frac{PL}{\pi d^2} \quad \frac{i^2 PL \times 4}{\pi d^2} = 0.5 C \times 40$$

$$\frac{i^2 PL}{\pi d^2} = 5C$$

Final result:


$$\frac{5 \times 60}{5\pi d^2} = 0.5 C \times 40$$

$$\frac{5 \times 60}{5} = 0.5 \times 40$$

3

PHYSICS

3. A transparent thin film of uniform thickness and refractive index $n_1 = 1.4$ is coated on the convex spherical surface of radius R at one end of a long solid glass cylinder of refractive index $n_2 = 1.5$, as shown in the figure. Rays of light parallel to the axis of the cylinder traversing through the film from air to glass get focused at distance f_1 from the film, while rays of light traversing from glass to air get focused at distance f_2 from the film. Then



- (A) $|f_1| = 3R$
- (B) $|f_1| = 2.8R$
- (C) $|f_2| = 2R$
- (D) $|f_2| = 1.4R$

4. A student is performing an experiment using a resonance column and a tuning fork of frequency 244 s^{-1} . He is told that the air in the tube has been replaced by another gas (assume that the column remains filled with the gas). If the minimum height at which resonance occurs is $(0.350 \pm 0.005) \text{ m}$, the gas in the tube is

(Useful information : $\sqrt{167RT} = 640 \text{ J}^{1/2} \text{ mole}^{-1/2}$; $\sqrt{140RT} = 590 \text{ J}^{1/2} \text{ mole}^{-1/2}$. The molar masses M in grams are given in the options. Take the values of $\sqrt{\frac{10}{M}}$ for each gas as given there.)

- (A) Neon ($M = 20, \sqrt{\frac{10}{20}} = \frac{7}{10}$)
- (B) Nitrogen ($M = 28, \sqrt{\frac{10}{28}} = \frac{3}{5}$)
- (C) Oxygen ($M = 32, \sqrt{\frac{10}{32}} = \frac{9}{16}$)
- (D) Argon ($M = 36, \sqrt{\frac{10}{36}} = \frac{17}{32}$)

Space for Rough Work

$\frac{1.4}{v} + \frac{1}{\infty} = \frac{0.4}{R}$ $\frac{1.4}{v} = \frac{0.4}{R}$ $\frac{1.4}{v} = \frac{0.4}{R}$ $\frac{1.4}{v} = \frac{0.4}{R}$ $\frac{1.4}{v} = \frac{0.4}{R}$ $\frac{1.4}{v} = \frac{0.4}{R}$ $\frac{1.4}{v} = \frac{0.4}{R}$ $\frac{1.4}{v} = \frac{0.4}{R}$ $\frac{1.4}{v} = \frac{0.4}{R}$ $\frac{1.4}{v} = \frac{0.4}{R}$

$\frac{1.5}{v} - \frac{1.4 \times 2}{7R} = \frac{0.1}{R}$ $\frac{1.5}{v} - \frac{0.4}{R} = \frac{0.1}{R}$ $\frac{1.5}{v} = \frac{0.5}{R}$ $v = 3R$

$\frac{1.4}{v} + \frac{1.5}{\infty} = -\frac{0.1}{R}$ $\frac{1.4}{v} = -\frac{0.1}{R}$ $v = -14R$ $1.4RT$

$\frac{1}{v} + \frac{1.4}{14R} = -\frac{0.4}{R}$ $\frac{1}{v} = -\frac{0.4}{R} - \frac{0.1}{R} = -\frac{0.5}{R}$ $\frac{R}{0.5} = 2R$

$0.355 = \frac{v}{2}$ $0.355 \times 244 = \sqrt{\frac{10RT}{M}}$

$\frac{10 \times 0.350 \times 244}{590} = \frac{1}{\sqrt{M}}$ $0.350 \times 244 = \sqrt{\frac{10RT}{M}}$ $\sqrt{\frac{10RT}{100M}}$


* 0

PHYSICS

PHYSICS


5. In the figure, a ladder of mass m is shown leaning against a wall. It is in static equilibrium making an angle θ with the horizontal floor. The coefficient of friction between the wall and the ladder is μ_1 and that between the floor and the ladder is μ_2 . The normal reaction of the wall on the ladder is N_1 and that of the floor is N_2 . If the ladder is about to slip, then

- (A) $\mu_1 = 0$ $\mu_2 \neq 0$ and $N_2 \tan \theta = \frac{mg}{2}$
- (B) $\mu_1 \neq 0$ $\mu_2 = 0$ and $N_1 \tan \theta = \frac{mg}{2}$
- ✓ (C) $\mu_1 \neq 0$ $\mu_2 \neq 0$ and $N_2 = \frac{mg}{1 + \mu_1 \mu_2}$
- (D) $\mu_1 = 0$ $\mu_2 \neq 0$ and $N_1 \tan \theta = \frac{mg}{2}$



6. Two ideal batteries of emf V_1 and V_2 and three resistances R_1 , R_2 and R_3 are connected as shown in the figure. The current in resistance R_2 would be zero if

- ✓ (A) $V_1 = V_2$ and $R_1 = R_2 = R_3$
- (B) $V_1 = V_2$ and $R_1 = 2R_2 = R_3$
- (C) $V_1 = 2V_2$ and $2R_1 = 2R_2 = R_3$
- ✓ (D) $2V_1 = V_2$ and $2R_1 = R_2 = R_3$



$$\sum \frac{V_1}{R_1} = \frac{V_2}{R_2}$$

$$V_1 + V_2 = (R_1 + R_2) i$$

$$\frac{V_1 + V_2}{R_1 + R_2} = i$$

Space for Rough Work

$$f_2 = N_1 \quad \mu_2 N_2 = N_1 \quad \mu_1 \mu_2 N_2 + N_2 = mg$$


$$\mu_1 N_1 + N_2 = mg \quad \frac{V_1}{R_1} = \frac{V_2}{R_3} \quad N_2 = \frac{mg}{1 + \mu_1 \mu_2}$$

$$N_2 \cos \theta = \frac{N_1 \mu_2 \sin \theta}{2} + \frac{\mu_2 N_2 \mu_1 \sin \theta}{2} + \frac{\mu_1 N_1 \mu_2 \sin \theta}{2}$$

$$N_2 \cos \theta = 2N_1 \sin \theta + \mu_1 N_1 \sin \theta$$

$$\frac{mg \cos \theta}{1 + \mu_1 \mu_2} = 2N_1 \sin \theta (2 + \mu_1)$$

$$\frac{mg}{2(1 + \mu_1 \mu_2)(2 + \mu_1)} = N_1 \tan \theta$$



$$V_2 = i R_3$$

$$2 \frac{V_1}{R_1} = \frac{V_2}{R_3} \quad 2R_3 = R_1 \quad V_1 + V_2 = i R_1 + (i - i) R_3$$

$$V_1 + V_2 - i R_1 = V_2$$


$$i R_1 + V_2 = i R_1 + i R_3 \quad V_1 = i R_1 \quad V_1 + V_2 = i R_1 + i R_3$$



9. One end of a taut string of length $3m$ along the x axis is fixed at $x = 0$. The speed of the waves in the string is 100 ms^{-1} . The other end of the string is vibrating in the y direction so that stationary waves are set up in the string. The possible waveform(s) of these stationary waves is(are)

- (A) $y(t) = A \sin \frac{\pi x}{6} \cos \frac{50\pi t}{3}$ (B) $y(t) = A \sin \frac{\pi x}{3} \cos \frac{100\pi t}{3}$
 (C) $y(t) = A \sin \frac{5\pi x}{6} \cos \frac{250\pi t}{3}$ (D) $y(t) = A \sin \frac{5\pi x}{2} \cos 250\pi t$

10. A parallel plate capacitor has a dielectric slab of dielectric constant K between its plates that covers $1/3$ of the area of its plates, as shown in the figure. The total capacitance of the capacitor is C while that of the portion with dielectric in between is C_1 . When the capacitor is charged, the plate area covered by the dielectric gets charge Q_1 and the rest of the area gets charge Q_2 . The electric field in the dielectric is E_1 and that in the other portion is E_2 . Choose the correct option/options, ignoring edge effects.



- (A) $\frac{E_1}{E_2} = 1$ (B) $\frac{E_1}{E_2} = \frac{1}{K}$ (C) $\frac{Q_1}{Q_2} = \frac{3}{K}$ (D) $\frac{C}{C_1} = \frac{2+K}{K}$

Space for Rough Work

Handwritten rough work for question 10:

$\frac{2\pi}{\lambda} = \frac{\pi}{6} \implies 12 = \lambda$ $\frac{2\pi}{\lambda} = \frac{\pi}{6} \implies \lambda = 12$ $\frac{2\pi}{\lambda} = \frac{\pi}{6} \implies \lambda = 12$ $\frac{2\pi}{\lambda} = \frac{\pi}{6} \implies \lambda = 12$ $\frac{2\pi}{\lambda} = \frac{\pi}{6} \implies \lambda = 12$ $\frac{2\pi}{\lambda} = \frac{\pi}{6} \implies \lambda = 12$

$\frac{5\pi}{6} = \frac{2\pi}{\lambda} \implies \lambda = \frac{12}{5}$ $\frac{5\pi}{6} = \frac{2\pi}{\lambda} \implies \lambda = \frac{12}{5}$ $\frac{5\pi}{6} = \frac{2\pi}{\lambda} \implies \lambda = \frac{12}{5}$ $\frac{5\pi}{6} = \frac{2\pi}{\lambda} \implies \lambda = \frac{12}{5}$

$C = \frac{2A\epsilon_0}{3d}$ $C_1 = \frac{A\epsilon_0 K}{3d}$ $C = \frac{2A\epsilon_0}{3d} + \frac{A\epsilon_0 K}{3d}$

$\frac{A\epsilon_0 K V}{3d} + \frac{2A\epsilon_0 V}{3d} = \frac{A\epsilon_0 K V}{3d} + \frac{2A\epsilon_0 V}{3d}$

$*0 \implies \frac{2A\epsilon_0 V}{3d} = \frac{A\epsilon_0 (K+2)V}{3d}$ $C_1 = \frac{A\epsilon_0 K}{3d}$ $Q_1 = \frac{A\epsilon_0 K V}{3d}$ $Q_2 = \frac{2A\epsilon_0 V}{3d}$

SECTION - 2 : (One Integer Value Correct Type)

This section contains 10 questions. Each question, when worked out will result in one integer from 0 to 9 (both inclusive).

11. A galvanometer gives full scale deflection with 0.006 A current. By connecting it to a 4990 Ω resistance, it can be converted into a voltmeter of range 0 - 30 V. If connected to a $\frac{2n}{249}$ Ω resistance, it becomes an ammeter of range 0 - 1.5 A. The value of n is 5
12. To find the distance d over which a signal can be seen clearly in foggy conditions, a railways engineer uses dimensional analysis and assumes that the distance depends on the mass density ρ of the fog, intensity (power/area) S of the light from the signal and its frequency f. The engineer finds that d is proportional to S^{1/n}. The value of n is 3

Space for Rough Work

$$\frac{30}{10} = 6 \times 10^{-3} (R + 4990)$$

$$5000 = R + 4990 \quad R = 10$$

$$6 \times 10^{-2} = (1.5 - 10) S$$

$$6 \times 10^{-2} = (1.5 - 0.6 \times 10^{-2}) S$$

$$\frac{6 \times 10^{-2}}{1.44} = S$$

kg × T S f

$$\frac{\text{kg}}{\text{L}} \times \frac{\text{J}}{\text{A S}}$$

$$\frac{\text{M}^1}{\text{L}^1 \text{T}} \times \frac{\text{M}^1 \text{L}^2 \text{T}^{-3}}{\text{L}^2}$$

$$\text{M}^2 \text{L}^{-1} \text{T}^{-4}$$

$$30 = 6 \times 10^{-3} (R_6 + 4990)$$

$$5000 = R_6 + 4990 \quad R_6 = 10$$

$$6 \times 10^{-3} \times 10 = (1.5 - 6 \times 10^{-3}) S$$


$$6 \times 10^{-2} = 1.494 S$$

$$\frac{60}{10} = S$$

$$6 = S$$



13. During Searle's experiment, zero of the Vernier scale lies between $3.20 \times 10^{-2} \text{ m}$ and $3.25 \times 10^{-2} \text{ m}$ of the main scale. The 20th division of the Vernier scale exactly coincides with one of the main scale divisions. When an additional load of 2 kg is applied to the wire, the zero of the Vernier scale still lies between $3.20 \times 10^{-2} \text{ m}$ and $3.25 \times 10^{-2} \text{ m}$ of the main scale but now the 45th division of Vernier scale coincides with one of the main scale divisions. The length of the thin metallic wire is 2 m and its cross-sectional area is $8 \times 10^{-7} \text{ m}^2$. The least count of the Vernier scale is $1.0 \times 10^{-5} \text{ m}$. The maximum percentage error in the Young's modulus of the wire is
14. A horizontal circular platform of radius 0.5 m and mass 0.45 kg is free to rotate about its axis. Two massless spring toy-guns, each carrying a steel ball of mass 0.05 kg are attached to the platform at a distance 0.25 m from the centre on its either sides along its diameter (see figure). Each gun simultaneously fires the balls horizontally and perpendicular to the diameter in opposite directions. After leaving the platform, the balls have horizontal speed of 9 ms^{-1} with respect to the ground. The rotational speed of the platform in rad s^{-1} after the balls leave the platform is



Space for Rough Work

$$Y = \frac{F \ell}{A \Delta l}$$

$$\frac{0.05 \times 10^{-2}}{20} \quad 20 \times 10^{-5}$$

$$\log Y =$$

$$45 \times 10^{-5}$$

$$32 \times 10^{-3} \quad 32.2 \times 10^{-3}$$

$$0.45 \times 10^{-3}$$

$$32.45 \times 10^{-3}$$

$$L \times [M^1 L^{-3}]^a [M^1 T^{-3}]^b [T^{-1}]^c$$

$$= M^{1+ab} L^{-3a} T^{-3b-c}$$

$$0.05 \times 2 \times 9 \times 0.25$$

$$= \frac{1}{2} \times 0.45 \times 0.25 \times \omega$$

$$-3a = 1$$

$$a = -\frac{1}{3}$$

$$-\frac{1}{3} + b = 0$$

$$b = \frac{1}{3}$$

$$-3b - c = 0$$

$$c = -1$$


$$\frac{0.01}{0.45 \times 4 \times 900} = \omega$$

$$\frac{0.45 \times 45}{5} = \omega$$




PHYSICS

15. A uniform circular disc of mass 1.5 kg and radius 0.5 m is initially at rest on a horizontal frictionless surface. Three forces of equal magnitude $F = 0.5 \text{ N}$ are applied simultaneously along the three sides of an equilateral triangle XYZ with its vertices on the perimeter of the disc (see figure). One second after applying the forces, the angular speed of the disc in rad s^{-1} is



~~cos 30~~ = $\frac{\sqrt{3}}{2} = \frac{x}{0.5}$
 $\frac{\sqrt{3}}{4} = x$
 $\sin 30 = \frac{1}{2} = \frac{x}{0.5}$

16. Consider an elliptically shaped rail PQ in the vertical plane with $OP = 3 \text{ m}$ and $OQ = 4 \text{ m}$. A block of mass 1 kg is pulled along the rail from P to Q with a force of 18 N, which is always parallel to line PQ (see the figure given). Assuming no frictional losses, the kinetic energy of the block when it reaches Q is $(n \times 10) \text{ Joules}$. The value of n is (take acceleration due to gravity = 10 ms^{-2})




~~work~~
 $18 = 9$
 $v^2 = \frac{2 \times 18 \times 5}{2}$
 $\frac{18}{90}$

Space for Rough Work

$\frac{3 \times \sqrt{3}}{4} \times \frac{1}{2} = \frac{1}{2} \times 1.5 \times \frac{1}{4} \alpha$
 $2\sqrt{3} = \alpha$
 $\frac{x^2}{9} + \frac{y^2}{16} = 1$
 $\frac{2x}{9} + \frac{2yy'}{16} = 0$
 $\frac{4y'}{16} = -\frac{x}{9}$
 $y' = -\frac{16x}{9y}$
 $3 \times \frac{1}{2} \times \frac{1}{4} = \frac{1}{2} \times 1.5 \times \frac{1}{4} \alpha$
 $2 = \alpha$
 $\omega = \alpha t$
 $R_1 = \frac{mV}{9 \times 300 I}$
 $R_2 = \frac{mV}{9 \times 900 I}$
 $\frac{mV}{39 \times 300 I}$



17. A thermodynamic system is taken from an initial state i with internal energy $U_i = 100\text{ J}$ to the final state f along two different paths iaf and ibf , as schematically shown in the figure. The work done by the system along the paths af , ib and bf are $W_{af} = 200\text{ J}$, $W_{ib} = 50\text{ J}$ and $W_{bf} = 100\text{ J}$ respectively. The heat supplied to the system along the path iaf , ib and bf are Q_{iaf} , Q_{ib} and Q_{bf} respectively. If the internal energy of the system in the state b is $U_b = 200\text{ J}$ and $Q_{iaf} = 500\text{ J}$, the ratio Q_{bf}/Q_{ib} is



Handwritten calculations for problem 17:

$$500 - 200 = Q_{ibf}$$

$$-150$$

$$300$$

$$150$$

$$450 = Q_{ib}$$

$$Q_{ib} = 100 + 50$$

18. Two parallel wires in the plane of the paper are distance X_0 apart. A point charge is moving with speed u between the wires in the same plane at a distance X_1 from one of the wires. When the wires carry current of magnitude I in the same direction, the radius of curvature of the path of the point charge is R_1 . In contrast, if the currents I in the two wires have directions opposite to each other, the radius of curvature of the path is R_2 . If $\frac{X_0}{X_1} = 3$, the value of $\frac{R_1}{R_2}$ is

Handwritten calculation for problem 18:

$$Q_{bf} = 450$$


$$150$$

$$300$$

Space for Rough Work

Handwritten equation:

$$\frac{\mu_0 I}{2\pi r_1} - \frac{\mu_0 I}{2\pi (X_0 - r_1)}$$



Handwritten equation:

$$\frac{X_0}{X_1} = 3$$

Handwritten equation:

$$\left(\frac{\mu_0 I}{2\pi r_1} - \frac{\mu_0 I}{2\pi (X_0 - r_1)} \right)$$

Handwritten equation:

$$\frac{mv}{q \frac{\mu_0 I}{2\pi r_1} - \mu_0}$$

Handwritten equation:

$$Q_{ib} = 100 + 50$$

Handwritten equation:

$$500 + 200 = Q_{ib} + W_{ib} + Q_{bf} + W_{bf}$$

Handwritten equation:

$$700 = Q_{ib} + Q_{bf} + 150$$

Handwritten equation:

$$Q_{ibf} = 550$$

Handwritten calculations for problem 18:

$$300$$

$$150$$

$$610$$

$$760$$

$$150$$

$$550$$

$$150$$

$$550$$


$$150$$

$$400$$




PHYSICS

19. Airplanes A and B are flying with constant velocity in the same vertical plane at angles 30° and 60° with respect to the horizontal respectively as shown in figure. The speed of A is $100\sqrt{3} \text{ ms}^{-1}$. At time $t = 0 \text{ s}$, an observer in A finds B at a distance of 500 m . This observer sees B moving with a constant velocity perpendicular to the line of motion of A. If at $t = t_0$, A just escapes being hit by B, t_0 in seconds is



20. A rocket is moving in a gravity free space with a constant acceleration of 2 ms^{-2} along $+x$ direction (see figure). The length of a chamber inside the rocket is 4 m . A ball is thrown from the left end of the chamber in $+x$ direction with a speed of 0.3 ms^{-1} relative to the rocket. At the same time, another ball is thrown in $-x$ direction with a speed of 0.2 ms^{-1} from its right end relative to the rocket. The time in seconds when the two balls hit each other is



Space for Rough Work

~~0.3~~ $v_{rel} = 0.3 + 0.2 = 0.5$ $s_{ru} = 40$

t $0.3 - 2t$ $0.2 + 2t$

$0.3 - \frac{4}{0.5} = \frac{40}{5} - 8$ $v_{BR} = v_B - v_R = 0.3 - 2t - 0.2 - 2t$

$A = 15t + \frac{1}{2} \times 2 \times t^2$
 $A = 15t + t^2$
 $B = 10t - \frac{1}{2} \times 2 \times t^2$
 $B = 10t - t^2$
 $20t = 20$
 $t = 1$



PART II : CHEMISTRY

SECTION – 1 : (One or More Than One Options Correct Type)

This section contains 10 multiple choice type questions. Each question has four choices (A), (B), (C) and (D) out of which ONE or MORE THAN ONE are correct.

21. Upon heating with Cu_2S , the reagent(s) that give copper metal is/are

- (A) CuFeS_2 (B) CuO
 (C) Cu_2O (D) CuSO_4

22. In a galvanic cell, the salt bridge

- (A) does not participate chemically in the cell reaction.
 (B) stops the diffusion of ions from one electrode to another.
 (C) is necessary for the occurrence of the cell reaction.
(D) ensures mixing of the two electrolytic solutions.

CHEMISTRY

Space for Rough Work

$$4 = 0.2t + t^2$$
$$t^2 + 0.2t - 4 = 0$$
$$t = \frac{-0.2 \pm \sqrt{0.04 + 16}}{2}$$
$$t = \frac{-0.2 \pm 4}{2}$$
$$= \frac{3.8}{2} = 1.9$$




23. Hydrogen bonding plays a central role in the following phenomena:

- (A) Ice floats in water.
- (B) Higher Lewis basicity of primary amines than tertiary amines in aqueous solutions.
- (C) Formic acid is more acidic than acetic acid.
- (D) Dimerisation of acetic acid in benzene.

24. The correct combination of names for isomeric alcohols with molecular formula $C_4H_{10}O$ is/are

- (A) *tert*-butanol and 2-methylpropan-2-ol
- (B) *tert*-butanol and 1, 1-dimethylethan-1-ol
- (C) *n*-butanol and butan-1-ol
- (D) isobutyl alcohol and 2-methylpropan-1-ol



25. The reactivity of compound **Z** with different halogens under appropriate conditions is given below :



The observed pattern of electrophilic substitution can be explained by

- (A) the steric effect of the halogen
- (B) the steric effect of the *tert*-butyl group
- (C) the electronic effect of the phenolic group
- (D) the electronic effect of the *tert*-butyl group

Space for Rough Work



26. The pair(s) of reagents that yield paramagnetic species is/are

- (A) Na and excess of NH_3
- (B) K and excess of O_2
- (C) Cu and dilute HNO_3
- (D) O_2 and 2-ethylantraquinol




27. For the reaction :





The correct statement(s) in the balanced equation is/are :

- (A) Stoichiometric coefficient of HSO_4^- is 6.
- (B) Iodide is oxidized.
- (C) Sulphur is reduced.
- (D) H_2O is one of the products.






Space for Rough Work




~~ClO₃⁻~~




28. In the reaction shown below, the major product(s) formed is/are



Space for Rough Work



29. An ideal gas in a thermally insulated vessel at internal pressure = P_1 , volume = V_1 and absolute temperature = T_1 expands irreversibly against zero external pressure, as shown in the diagram. The final internal pressure, volume and absolute temperature of the gas are P_2 , V_2 and T_2 , respectively. For this expansion,



- (A) $q = 0$
 (B) $T_2 = T_1$
 (C) $P_2V_2 = P_1V_1$
 (D) $P_2V_2^\gamma = P_1V_1^\gamma$

30. The correct statement(s) for orthoboric acid is/are

- (A) It behaves as a weak acid in water due to self ionization.
 (B) Acidity of its aqueous solution increases upon addition of ethylene glycol.
 (C) It has a three dimensional structure due to hydrogen bonding.
 (D) It is a weak electrolyte in water.


Space for Rough Work



SECTION – 2 : (One Integer Value Correct Type)


This section contains 10 questions. Each question, when worked out will result in one integer from 0 to 9 (both inclusive).

31. The total number of distinct naturally occurring amino acids obtained by complete acidic hydrolysis of the peptide shown below is



32. MX_2 dissociates into M^{2+} and X^- ions in an aqueous solution, with a degree of dissociation (α) of 0.5. The ratio of the observed depression of freezing point of the aqueous solution to the value of the depression of freezing point in the absence of ionic dissociation is 2
33. If the value of Avogadro number is $6.023 \times 10^{23} \text{ mol}^{-1}$ and the value of Boltzmann constant is $1.380 \times 10^{-23} \text{ J K}^{-1}$, then the number of significant digits in the calculated value of the universal gas constant is 3
34. A compound H_2X with molar weight of 80 g is dissolved in a solvent having density of 0.4 g ml^{-1} . Assuming no change in volume upon dissolution, the **molality** of a 3.2 molar solution is 2
35. In an atom, the total number of electrons having quantum numbers $n = 4$, $|m_l| = 1$ and $m_s = -1/2$ is 6


Space for Rough Work



$$\frac{R}{N_A} = K$$

*0

$$R = 6.023 \times 10^{23} \Delta T_f = 2 K_f m$$



18

$$MX_2 \rightleftharpoons M^{2+} + 2X^-$$

C	0	0
C	0.5C	0.5C
	0.5C	0.5C

$$\frac{2R}{2} = i$$

$$2 = 1$$




36. Consider the following list of reagents :
 Acidified $K_2Cr_2O_7$, alkaline $KMnO_4$, $CuSO_4$, H_2O_2 , Cl_2 , O_3 , $FeCl_3$, HNO_3 and $Na_2S_2O_3$.
 The total number of reagents that can oxidise aqueous iodide to iodine is 8

37. A list of species having the formula XZ_4 is given below.
 XeF_4 , SF_4 , SiF_4 , BF_4^- , BrF_4^- , $[Cu(NH_3)_4]^{2+}$, $[FeCl_4]^{2-}$, $[CoCl_4]^{2-}$ and $[PtCl_4]^{2-}$.
 Defining shape on the basis of the location of X and Z atoms, the total number of species having a square planar shape is

38. Consider all possible isomeric ketones, including stereoisomers of MW = 100. All these isomers are independently reacted with $NaBH_4$ (NOTE: stereoisomers are also reacted separately). The total number of ketones that give a racemic product(s) is/are

39. The total number(s) of stable conformers with **non-zero** dipole moment for the following compound is (are)



40. Among PbS , CuS , HgS , MnS , Ag_2S , NiS , CoS , Bi_2S_3 and SnS_2 , the total number of **BLACK** coloured sulphides is 7

CHEMISTRY

Space for Rough Work

Handwritten calculations and diagrams for rough work:

- $3.2 \text{ mol} \equiv 1 \text{ L}$
- $3.2 \times 80 \text{ g} \equiv 1 \text{ L } H_2X$
- $0.4 \text{ g sol} \equiv 1 \text{ ml}$
- $1000 \text{ ml} \equiv 400 \text{ g}$
- $1 \text{ L} \equiv 400 \text{ g}$
- $3.2 \times 1000 = 3200$
- 144
- 1600
- $3200 + 144 = 3344$
- $3344 / 72 = 46.44$
- 2.2
- $72 \overline{) 1608}$
- 144
- 160
- 32
- 8
- 4
- 256
- $39, 10$
- 400
- 256
- 164
- 5
- 0.3
- 72
- 19
- $1/2(2+4) = 3$
- $1/2(5+4-1) = 4$
- $1/2(2+4-1) = 2.5$
- 72
- 12
- 5
- 88
- 2.5
- 87
- 48
- 16
- 101
- 8
- $2CO_2$
- $CH_3 - CH_2 - CH_2 - CH_2 - CH_2 - CH_3$

PART III : MATHEMATICS

SECTION – 1 : (One or More Than One Options Correct Type)

This section contains 10 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which ONE or MORE THAN ONE are correct.

41. For every pair of continuous functions $f, g: [0, 1] \rightarrow \mathbb{R}$ such that

$$\max \{f(x): x \in [0, 1]\} = \max \{g(x): x \in [0, 1]\},$$

the correct statement(s) is(are) :

- (A) $(f'(c))^2 + 3f(c) = (g'(c))^2 + 3g(c)$ for some $c \in [0, 1]$
- (B) $(f(c))^2 + f(c) = (g(c))^2 + 3g(c)$ for some $c \in [0, 1]$
- (C) $(f(c))^2 + 3f(c) = (g(c))^2 + g(c)$ for some $c \in [0, 1]$
- (D) $(f(c))^2 = (g(c))^2$ for some $c \in [0, 1]$

42. A circle S passes through the point $(0, 1)$ and is orthogonal to the circles $(x - 1)^2 + y^2 = 16$ and $x^2 + y^2 = 1$. Then

- (A) radius of S is 8
- (B) radius of S is 7
- (C) centre of S is $(-7, 1)$
- (D) centre of S is $(-8, 1)$

Space for Rough Work

$$g_1 = \sqrt{(h-1)^2 + k^2}$$

$$g_2 = \sqrt{h^2 + (k-1)^2}$$

$$\sqrt{(h-1)^2 + k^2} + 16 = \sqrt{h^2 + (k-1)^2}$$

$$h^2 + k^2 + 1 - 2k + 16 = h^2 + k^2 - 2k + 1$$

$$2h - 2k + 16 = 0 \quad h - k + 8 = 0$$

$$h^2 + (k-1)^2 + 1 = h^2 + k^2$$

$$k^2 + 1 - 2k + 1 = k^2$$

$$2 - 2k = 0 \quad k = 1$$

$$h - 1 + 8 = 0$$

$$h = -7$$

$$\vec{a} \cdot \vec{b} = (\vec{b} \cdot \vec{c}) \cdot (\vec{c} \cdot \vec{a})$$

$$(\vec{a} \cdot \vec{b}) \cdot (\vec{c} \cdot \vec{c}) = (\vec{b} \cdot \vec{c}) \cdot (\vec{c} \cdot \vec{a})$$

$$\vec{a} = \lambda (\vec{c} \times (\vec{b} \times \vec{c}))$$


$$= \lambda (\vec{c} \cdot \vec{c}) \vec{b} - (\vec{c} \cdot \vec{b}) \vec{c}$$

$$= \lambda (\vec{b} - \vec{c})$$

$$\vec{b} = \mu (\vec{c} \times (\vec{a} \times \vec{c}))$$

$$= \mu (\vec{c} \cdot \vec{c}) \vec{a} - (\vec{c} \cdot \vec{a}) \vec{c}$$

$$= \mu (\vec{a} - \vec{c})$$



MATHEMATICS

43. From a point $P(\lambda, \lambda, \lambda)$, perpendiculars PQ and PR are drawn respectively on the lines $y = x, z = 1$ and $y = -x, z = -1$. If P is such that $\angle QPR$ is a right angle, then the possible value(s) of λ is(are)

- (A) $\sqrt{2}$ (B) 1 (C) -1 (D) $-\sqrt{2}$

44. Let \vec{x}, \vec{y} and \vec{z} be three vectors each of magnitude $\sqrt{2}$ and the angle between each pair of them is $\frac{\pi}{3}$. If \vec{a} is a nonzero vector perpendicular to \vec{x} and $\vec{y} \times \vec{z}$ and \vec{b} is a nonzero vector perpendicular to \vec{y} and $\vec{z} \times \vec{x}$, then

- (A) $\vec{b} = (\vec{b} \cdot \vec{z})(\vec{z} - \vec{x})$ (B) $\vec{a} = (\vec{a} \cdot \vec{y})(\vec{y} - \vec{z})$
 (C) $\vec{a} \cdot \vec{b} = -(\vec{a} \cdot \vec{y})(\vec{b} \cdot \vec{z})$ (D) $\vec{a} = (\vec{a} \cdot \vec{y})(\vec{z} - \vec{y})$

45. Let $f: (-\frac{\pi}{2}, \frac{\pi}{2}) \rightarrow \mathbb{R}$ be given by

$f(x) = (\log(\sec x + \tan x))^3$

Then

- (A) $f(x)$ is an odd function (B) $f(x)$ is a one-one function
 (C) $f(x)$ is an onto function (D) $f(x)$ is an even function

Space for Rough Work

MATHEMATICS

$x = y = 0, z = 1$
 $\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 0 \\ 0 & 0 & 1 \end{vmatrix} \Rightarrow 1(\hat{i} - \hat{j})$
 $-\hat{i} - \hat{j} = \vec{b}_1$
 $\frac{x-1}{-1} = \frac{y-1}{-1} = \frac{z-1}{0} = u$
 $Q(-u+1, -u+1, +1)$
 $\vec{OP} = (\lambda - u + 1)\hat{i} + (\lambda + u - 1)\hat{j} + (\lambda - 1)\hat{k}$
 $-\lambda + u + 1 - \lambda - u + 1 = 0$
 $\Rightarrow 2\lambda = 2 \Rightarrow \lambda = 1$
 $\vec{a} \cdot \vec{x} = 0$
 $\vec{a} \cdot (\vec{y} \times \vec{z}) = 0$
 $\vec{a} \cdot \vec{z} = 0$

$x + y, z + 1 = 0 \Rightarrow -1, -1, 1$
 $\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix}$
 $1(\hat{i} - \hat{j}) = \vec{b}_2$
 $\frac{x+1}{1} = \frac{y-1}{-1} = \frac{z+1}{0} = \alpha$
 $R(\alpha - 1, -\alpha + 1, -1)$
 $\vec{PR} = (\lambda - \alpha + 1)\hat{i} + (\lambda + \alpha - 1)\hat{j} + (\lambda - 1)\hat{k}$
 $\vec{b} \cdot \vec{y} = 0$
 $\vec{b} \cdot (\vec{z} \times \vec{x}) = 0$
 $\vec{b} \cdot \vec{x} = 0$
 $\lambda - \alpha + 1 - \lambda - \alpha + 1 = 0$
 $\Rightarrow 2\alpha = 2 \Rightarrow \alpha = 1$

$\frac{1 + \sin x}{\cos x}$
 $1 - \cos(\frac{\pi}{2} + x)$
 $\sin(\frac{\pi}{2} + x)$

462
140
5047

MATHEMATICS

46. Let $f: [a, b] \rightarrow [1, \infty)$ be a continuous function and let $g: \mathbb{R} \rightarrow \mathbb{R}$ be defined as

$$g(x) = \begin{cases} 0 & \text{if } x < a, \\ \int_a^x f(t) dt & \text{if } a \leq x \leq b, \\ \int_a^b f(t) dt & \text{if } x > b. \end{cases}$$

Then

- (A) $g(x)$ is continuous but not differentiable at a
- (B) $g(x)$ is differentiable on \mathbb{R}
- (C) $g(x)$ is continuous but not differentiable at b
- (D) $g(x)$ is continuous and differentiable at either a or b but not both

$f'(a^+) = \frac{f(a)}{0}$

47. Let $f: (0, \infty) \rightarrow \mathbb{R}$ be given by

$$f(x) = \int_{\frac{1}{x}}^x e^{-(t+\frac{1}{t})} \frac{dt}{t}.$$

Then

- (A) $f(x)$ is monotonically increasing on $[1, \infty)$
- (B) $f(x)$ is monotonically decreasing on $(0, 1)$
- (C) $f(x) + f(\frac{1}{x}) = 0$, for all $x \in (0, \infty)$
- (D) $f(2^x)$ is an odd function of x on \mathbb{R}

Space for Rough Work

$\lim_{x \rightarrow a^-} g(x) = 0$ $\lim_{x \rightarrow a^+} \int_0^x f(t) dt = 0$
 $\lim_{x \rightarrow b^-} \int_a^b f(t) dt = \int_a^b f(t) dt$ $-\int_a^b \frac{1}{t} e^{-(\frac{1}{t}+t)} \frac{1}{t^2}$
 $f(x) = \int_{\frac{1}{x}}^x \frac{e^{-(t+\frac{1}{t})}}{t} dt$ $f(x) = \frac{e^{-(x+\frac{1}{x})}}{x} + \frac{e^{-(\frac{1}{x}+x)}}{x^2}$
 $f'(x) = \frac{e^{-(x+\frac{1}{x})}}{x^2} - \frac{e^{-(\frac{1}{x}+x)}}{x^3}$
 $f'(x) \rightarrow 0$
 $\int_{\frac{1}{x}}^x \frac{e^{-(t+\frac{1}{t})}}{t} dt = \int_x^{\frac{1}{x}} \frac{e^{-(t+\frac{1}{t})}}{t} dt$
 $t = \frac{1}{u} \quad dt = -\frac{1}{u^2} du$



48. Let M be a 2×2 symmetric matrix with integer entries. Then M is invertible if
- (A) the first column of M is the transpose of the second row of M
 - (B) the second row of M is the transpose of the first column of M
 - (C) M is a diagonal matrix with nonzero entries in the main diagonal
 - (D) the product of entries in the main diagonal of M is not the square of an integer

49. Let $a \in \mathbb{R}$ and let $f: \mathbb{R} \rightarrow \mathbb{R}$ be given by

$$f(x) = x^5 - 5x + a.$$

Then

- (A) $f(x)$ has three real roots if $a > 4$
 - (B) $f(x)$ has only one real root if $a > 4$
 - (C) $f(x)$ has three real roots if $a < -4$
 - (D) $f(x)$ has three real roots if $-4 < a < 4$
50. Let M and N be two 3×3 matrices such that $MN = NM$. Further, if $M \neq N^2$ and $M^2 = N^4$, then
- (A) determinant of $(M^2 + MN^2)$ is 0
 - (B) there is a 3×3 non-zero matrix U such that $(M^2 + MN^2)U$ is the zero matrix
 - (C) determinant of $(M^2 + MN^2) \geq 1$
 - (D) for a 3×3 matrix U , if $(M^2 + MN^2)U$ equals the zero matrix then U is the zero matrix

Space for Rough Work

$$[\quad]$$

$$MN = NM$$

$$M^2 = N^4$$

$$N^4 + MN^2$$

$$(N^2 + M)N^2$$

$$\det M^2 = \det N^4$$

$$(\det M)^2 = (\det N)^4$$

$$\Rightarrow \det M = \det N$$

$$M^2 + MN^2$$

$$M^2 + NMN$$



$$2 \begin{vmatrix} 1 & 1 & 1 \\ \frac{1}{2} & 1 & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & 1 \end{vmatrix} = 2 \begin{vmatrix} 1 & 0 & 0 \\ \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \end{vmatrix} = 2 + \frac{1}{2} = \frac{5}{2}$$

SECTION - 2 : (One Integer Value Correct Type)

This section contains 10 questions. Each question, when worked out will result in one integer from 0 to 9 (both inclusive).

51. For a point P in the plane, let $d_1(P)$ and $d_2(P)$ be the distances of the point P from the lines $x - y = 0$ and $x + y = 0$ respectively. The area of the region R consisting of all points P lying in the first quadrant of the plane and satisfying $2 \leq d_1(P) + d_2(P) \leq 4$, is **6** ✓

52. Let \vec{a} , \vec{b} , and \vec{c} be three non-coplanar unit vectors such that the angle between every pair of them is $\frac{\pi}{3}$. If $\vec{a} \times \vec{b} + \vec{b} \times \vec{c} = p\vec{a} + q\vec{b} + r\vec{c}$, where p, q and r are scalars, then the value of $\frac{p^2 + 2q^2 + r^2}{q^2}$ is **7** (X)

53. The largest value of the non-negative integer a for which

$$\lim_{x \rightarrow 1} \left\{ \frac{-ax + \sin(x-1) + a}{x + \sin(x-1) - 1} \right\}^{\frac{1-x}{1-\sqrt{x}}} = \frac{1}{4}$$

is **0**


54. Let $f: [0, 4\pi] \rightarrow [0, \pi]$ be defined by $f(x) = \cos^{-1}(\cos x)$. The number of points $x \in [0, 4\pi]$ satisfying the equation

$$f(x) = \frac{10-x}{10}$$

is **3**

Space for Rough Work

Handwritten rough work for question 54. It includes a coordinate system with points $(0, 2\sqrt{2})$, $(\sqrt{2}, 2\sqrt{2})$, $(2\sqrt{2}, 0)$, and $(2\sqrt{2}, 2\sqrt{2})$ marked. A square is drawn with vertices at $(0, \sqrt{2})$, $(\sqrt{2}, \sqrt{2})$, $(\sqrt{2}, 0)$, and $(0, 0)$. The origin is marked with $*0$. The number 24 is written. The equation $2 \leq \frac{|h-k| + |h+k|}{\sqrt{2}} \leq 4$ is derived, leading to $|h-k| + |h+k| \leq 4\sqrt{2}$. Further steps show $h \leq 2\sqrt{2}$ and $k \leq 2\sqrt{2}$. A barcode is present at the bottom right of the rough work area.



MATHEMATICS

55. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ be respectively given by $f(x) = |x| + 1$ and $g(x) = x^2 + 1$. Define $h: \mathbb{R} \rightarrow \mathbb{R}$ by

$$h(x) = \begin{cases} \max \{f(x), g(x)\} & \text{if } x \leq 0, \\ \min \{f(x), g(x)\} & \text{if } x > 0. \end{cases}$$

The number of points at which $h(x)$ is not differentiable is

3 ✓

56. Let $n_1 < n_2 < n_3 < n_4 < n_5$ be positive integers such that $n_1 + n_2 + n_3 + n_4 + n_5 = 20$. Then the number of such distinct arrangements $(n_1, n_2, n_3, n_4, n_5)$ is

1, 2, 3, 4, 5

57. Let $n \geq 2$ be an integer. Take n distinct points on a circle and join each pair of points by a line segment. Colour the line segment joining every pair of adjacent points by blue and the rest by red. If the number of red and blue line segments are equal, then the value of n is

5 ✓


Space for Rough Work

$nC - n = n$

$20 + 5 - 1$

$\frac{n(n-1)}{2} = 2n, n = 5$

1, 3, 4, 5



$y - 1 = |x|$

$y - 1 = x^2$

MATHEMATICS

$\lim_{x \rightarrow 1} \frac{-ax + \sin(x-1) + a}{x + \sin(x-1)} = \frac{1}{2} \quad y = -x + 1$

$\frac{-a + \cos(x-1)}{1 + \cos(x-1)} = \frac{-a+1}{1+1} = \frac{1}{2}$

$\cos^{-1} \cos x = \frac{10-x}{10} \quad \frac{-a+1}{2} = \frac{1}{2}$

$\cos x = \cos \frac{10-x}{10}$

$x + 10 = 40x \quad 10x = 10 - x$

$x = 2m\pi + \frac{10-x}{10} \quad x = \frac{10}{11}$

$\frac{10x + 10 - x}{11} = 4\pi \quad x + \frac{10-x}{10} = 4\pi$

Handwritten notes and calculations including $y = -x + 1$ for $x < 0$ and $y = x^2 + 1$ for $x > 0$, and various numerical manipulations.



$$2(y-x^5)(y'-5x^4) = (1+x^2)^2 + 2x(1+x^2)2x$$

$$2 \times 2 (y'-5) = \underline{4+8}$$

MATHEMATICS

58. The slope of the tangent to the curve $(y - x^5)^2 = x(1 + x^2)^2$ at the point (1, 3) is

8

59. Let a, b, c be positive integers such that $\frac{b}{a}$ is an integer. If a, b, c are in geometric progression and the arithmetic mean of a, b, c is $b + 2$, then the value of

$$\frac{a^2 + a - 14}{a + 1}$$

is

4

$$\frac{a+b+c}{3} = b+2$$

$$a+c = 2b+6$$

$$b^2 = ac$$

$$\frac{a(a+1)-14}{a+1}$$

1

14/21

60. The value of

$$\int_0^1 4x^3 \left\{ \frac{d^2}{dx^2} (1-x^2)^5 \right\} dx$$

2

is $\frac{a \pm \sqrt{a}}{a}$

$$1 \pm \sqrt{\frac{6}{a}}$$

Space for Rough Work

$a = 6$

$a = 6$

$$\frac{d^2}{dx^2} (1-x^2)^5 = -2 \times 5 (1-x^2)^4 \times 2x$$

$$= -10x(1-x^2)^4$$

24 ± 12

$$2(y-x^5)(y'-5x^4) = (1+x^2)^2 + 2x(1+x^2)2x$$

$$2(2)(y'-5) = 4 + 4 \times 2$$

$$4(y'-5) = 12$$

$$y' = 8$$

$$-40x(1-x^2)^3(-2x) - 10(1-x^2)^4$$

$$80x^2(1-x^2)^3 - 10(1-x^2)^4$$

$$4x^3 \cdot 10(1-x^2)^3(8x^2-1)$$

$$2(2)(y'-5) =$$

$$4 + 4 \times 2$$

$$4(y'-5) = 12$$

$$y' = 8$$

$$4x^3 \frac{d}{dx} (1-x^2)^5 \Big|_0^1$$

$$- \int_0^1 12x^2 \frac{d}{dx} (1-x^2)^5 dx$$

$$-12x^2 (1-x^2)^5 \Big|_0^1 + \int_0^1 24x(1-x^2)^5 dx$$

*0

$$12 \int_0^1 t^5 dt = 2$$

$$a+ar+ar^2 = 3ar+6$$

$$a+ar^2 = 2ar+6$$

$$ar^2 - 2ar + a - 6 = 0$$

$$r = \frac{2a \pm \sqrt{4a^2 - 4a(a-6)}}{2a}$$

$$\frac{a+ar+ar^2}{3} = ar+2$$

$$a+ar^2 = 2ar+6$$

$$2a \pm \sqrt{24a}$$

$$a \pm \sqrt{6a}$$

12

$$1-x^2 = t$$

$$-2x dx = dt$$

$$ar^2 - 2ar + a - 6 = 0$$

$$r = \frac{2a \pm \sqrt{4a^2 - 4a(a-6)}}{2a}$$